

Abstract Submitted to the
International Conference on Strongly Correlated Electron Systems
University of Michigan, Ann Arbor
August 6-10, 2001

Spin splitting of the conduction band of the ferromagnetic semiconductor EuO

P.G. Steeneken¹, L.H. Tjeng¹, I. Elfimov¹, G.A. Sawatzky¹, G. Ghiringhelli², N.B. Brookes³, D.-J. Huang⁴

¹ *Solid State Physics Laboratory, University of Groningen, Nijenborgh 4, 9747 AG Groningen, the Netherlands*

² *Dipartimento di Fisica, Politecnico di Milano, piazza Leonardo da Vinci 32, 20133 Milano, Italy*

³ *European Synchrotron Radiation Facility, B.P. 220, 38043 Grenoble Cedex, France*

⁴ *Synchrotron Radiation Research Center, No. 1 R&D VI, Hsinchu Science-based Industrial Park, Hsinchu 30077, Taiwan*

EuO is a semiconductor with a band gap of 1.2 eV and is one of the rare ferromagnetic oxides. It becomes metallic below its Curie temperature of 69 K via a spectacular metal-insulator transition: the resistivity drops by as much as 8 orders of magnitude. Moreover EuO shows giant Kerr rotations and colossal magnetoresistance which is much more extreme than in the manganates.

We have applied a new combination of spin-resolved photoemission, resonant-Auger and x-ray absorption measurements to determine the spin-polarized occupied and unoccupied density of states in EuO. These experiments revealed large changes over a wide energy range in the unoccupied density of states when varying the temperature across T_c . Moreover we clearly observed a splitting of 0.5 eV between the spin-up and spin-down conduction band. These results show that electron doped EuO in the ferromagnetic state will have charge carriers with a 100% spin-polarization.